



PREVALENCE of URINARY BLADDER cancer in SWITZERLAND

Figure 1. Observed and projected number of patients having been diagnosed with malignant urinary bladder cancer by index year and years since diagnosis.



**Table 1. Malignant urinary bladder cancer (ICD-10 C67).
Limited-duration prevalence in Switzerland by sex, age and year.**

		Limited-duration prevalence ²															
		Men								Women							
		2000		2005		2010		2015		2000		2005		2010		2015	
Age ³	Time since diagnosis [years]	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹
0-49	up to 2	33	1.3	27	1.1	43	1.7	43	1.7	7	0.3	6	0.3	17	0.7	9	0.3
	up to 5	69	2.8	58	2.3	73	2.9	69	2.7	14	0.6	28	1.2	38	1.5	24	1.0
	up to 10	112	4.6	75	3.0	105	4.1	93	3.6	23	1.0	34	1.4	53	2.1	38	1.5
	2 to 5	36	1.5	31	1.2	30	1.2	26	1.0	6	0.3	22	0.9	21	0.9	16	0.6
	5 to 10	43	1.8	17	0.7	33	1.3	24	0.9	9	0.4	6	0.3	15	0.6	14	0.6
50-59	up to 2	162	35	183	38	135	26	130	21	39	8	34	7	31	6	26	4
	up to 5	312	67	299	61	236	44	237	39	68	15	51	11	71	14	62	10
	up to 10	461	98	392	80	321	60	311	51	107	23	75	15	98	19	90	15
	2 to 5	150	32	115	24	100	19	107	18	29	6	17	4	40	8	36	6
	5 to 10	149	32	94	19	85	16	74	12	39	8	24	5	27	5	28	5
60-69	up to 2	304	96	348	96	325	78	376	84	87	25	84	22	80	18	78	17
	up to 5	578	183	671	185	631	151	705	158	170	48	150	39	171	39	172	37
	up to 10	946	299	969	267	983	235	1041	234	290	83	226	58	247	56	260	57
	2 to 5	274	87	323	89	306	73	329	74	83	24	66	17	91	21	94	20
	5 to 10	367	116	298	82	352	84	336	75	120	34	76	20	76	17	88	19
70-79	up to 2	432	206	451	198	418	167	442	151	99	34	119	39	63	20	78	23
	up to 5	920	439	836	367	882	352	954	326	249	85	216	72	143	46	166	48
	up to 10	1633	778	1367	599	1368	546	1425	487	421	144	340	113	268	86	272	79
	2 to 5	488	233	385	169	464	185	512	175	150	51	98	32	80	26	88	26
	5 to 10	713	340	531	233	486	194	470	161	172	59	124	41	125	40	106	31
80+	up to 2	207	215	288	254	302	230	315	207	113	57	113	50	119	48	112	43
	up to 5	420	436	546	482	599	456	621	409	194	98	214	96	232	94	211	80
	up to 10	801	832	879	776	978	745	1011	666	358	180	343	154	363	148	331	126
	2 to 5	213	221	259	228	298	227	307	202	81	41	102	46	113	46	99	38
	5 to 10	381	396	333	294	378	288	390	257	164	82	129	58	131	53	120	45
All ages	up to 2	1137	32	1297	35	1223	32	1305	32	345	9	356	9	311	8	302	7
	up to 5	2299	65	2410	66	2421	62	2585	63	695	19	660	17	655	16	635	15
	up to 10	3952	112	3682	101	3755	97	3881	95	1199	33	1019	27	1029	26	991	24
	2 to 5	1161	33	1113	30	1198	31	1281	31	349	9	304	8	345	9	333	8
	5 to 10	1654	47	1273	35	1335	34	1295	32	505	14	360	9	374	9	356	9

¹ Proportion per 100'000

² Estimated with Counting/Projection-Method (see chapter B and C in document "Prevalence estimation methods")

³ Age at index date

**Table 2. Malignant urinary bladder cancer (ICD-10 C67).
Complete prevalence in Switzerland by sex and year.**

		Complete prevalence ²															
		Men								Women							
		2000		2005		2010		2015		2000		2005		2010		2015	
Age ³	Time since diagnosis [years]	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹
All ages	more than 10	1217	35	1472	40	1319	34	1553	38	498	14	660	17	568	14	668	16
	unlimited	5169	147	5154	141	5074	131	5434	133	1697	46	1680	44	1597	40	1658	40

¹ Proportion per 100'000

² Estimated with Age-Period-Cohort model PIAMOD (see chapter D in document "Prevalence estimation methods")

³ Age at index date

May 2014

Contact: Matthias Lorez, PhD MAS
Foundation National Institute for Cancer Epidemiology and Registration (NICER)
c/o Universität Zürich
Seilergraben 49
CH-8001 Zürich
ml@nicer.org