


PREVALENCE of cancer of COLON and RECTUM in SWITZERLAND

Figure 1. Observed and projected number of patients having been diagnosed with malignant cancer of colon and rectum by index year and years since diagnosis.



**Table 1. Malignant cancer of colon and rectum (ICD-10 C18-20).
Limited-duration prevalence in Switzerland by sex, age and year.**

		Limited-duration prevalence ²															
		Men								Women							
		2000		2005		2010		2015		2000		2005		2010		2015	
Age ³	Time since diagnosis [years]	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹
0-49	up to 2	131	5	185	8	189	7	178	7	171	7	187	8	181	7	177	7
	up to 5	247	10	368	15	387	15	351	14	329	14	328	14	409	17	383	15
	up to 10	370	15	473	19	621	24	531	21	448	19	471	20	584	24	530	21
	2 to 5	116	5	183	7	197	8	174	7	158	7	141	6	228	9	207	8
	5 to 10	122	5	105	4	235	9	180	7	118	5	143	6	175	7	146	6
50-59	up to 2	528	113	484	99	498	94	487	80	327	70	382	78	372	71	397	67
	up to 5	942	201	928	190	971	183	971	161	652	140	753	154	778	148	789	134
	up to 10	1251	267	1274	261	1366	257	1374	227	889	190	1078	221	1113	212	1131	192
	2 to 5	414	88	444	91	474	89	484	80	325	70	371	76	405	77	391	66
	5 to 10	309	66	346	71	395	74	403	67	236	51	325	67	335	64	342	58
60-69	up to 2	749	237	976	269	1084	259	1153	259	511	146	622	160	575	131	584	127
	up to 5	1534	486	2000	551	2224	531	2382	535	1051	300	1259	325	1271	290	1287	280
	up to 10	2231	706	2929	807	3394	811	3599	808	1651	471	1909	492	2108	481	2144	466
	2 to 5	784	248	1024	282	1139	272	1229	276	541	154	637	164	696	159	703	153
	5 to 10	697	221	929	256	1170	279	1217	273	600	171	650	168	837	191	857	186
70-79	up to 2	994	473	1109	486	1077	429	1221	417	733	250	835	278	771	248	811	236
	up to 5	1989	947	2348	1030	2347	936	2586	883	1431	488	1750	583	1592	512	1642	479
	up to 10	2996	1428	3519	1544	3808	1519	4107	1403	2357	803	2628	875	2649	852	2662	776
	2 to 5	995	474	1240	544	1270	507	1365	466	699	238	915	305	821	264	831	242
	5 to 10	1008	480	1170	513	1461	583	1521	520	926	316	878	292	1057	340	1020	297
80+	up to 2	463	480	629	555	644	491	664	437	619	312	777	348	797	325	783	298
	up to 5	1023	1063	1368	1208	1468	1118	1513	997	1315	663	1536	689	1760	717	1733	659
	up to 10	1764	1833	2269	2003	2655	2022	2767	1823	2251	1135	2555	1146	3071	1251	3020	1148
	2 to 5	560	582	739	652	824	627	850	560	696	351	759	340	963	392	951	361
	5 to 10	742	770	901	796	1186	904	1253	826	936	472	1018	457	1311	534	1287	489
All ages	up to 2	2865	81	3383	93	3493	90	3702	91	2360	64	2803	74	2696	68	2751	66
	up to 5	5734	163	7013	192	7397	191	7803	192	4779	130	5626	148	5810	145	5834	140
	up to 10	8612	245	10464	286	11843	306	12377	304	7596	206	8640	227	9524	239	9486	228
	2 to 5	2869	82	3629	99	3904	101	4101	101	2419	66	2823	74	3114	78	3083	74
	5 to 10	2878	82	3452	94	4447	115	4574	112	2816	76	3014	79	3715	93	3652	88

¹ Proportion per 100'000

² Estimated with Counting/Projection-Method (see chapter B and C in document "Prevalence estimation methods")

³ Age at index date

**Table 2. Malignant cancer of colon and rectum (ICD-10 C18-20).
Complete prevalence in Switzerland by sex and year.**

		Complete prevalence ²															
		Men								Women							
		2000		2005		2010		2015		2000		2005		2010		2015	
Age ³	Time since diagnosis [years]	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹	N	Prop ¹
All ages	more than 10	2810	80	3214	88	4109	106	6449	158	4026	109	4368	115	4839	121	6878	165
	unlimited	11422	325	13678	374	15952	412	18826	462	11622	315	13008	342	14364	360	16365	393

¹ Proportion per 100'000

² Estimated with Age-Period-Cohort model PIAMOD (see chapter D in document "Prevalence estimation methods")

³ Age at index date

May 2014

Contact: Matthias Lorez, PhD MAS
Foundation National Institute for Cancer Epidemiology and Registration (NICER)
c/o Universität Zürich
Seilergraben 49
CH-8001 Zürich
ml@nicer.org